

Lor et t e Communiqué

Published monthly by volunteers

Please e-mail up-coming events to news1@mts.net or news2@mts.net or phone 878-2353 or 878-3547

It's better to sleep on your decisions than to stay awake over what you have done.


Lorette Skating Club is holding an ice show At the Hop Fun from the '50s Lorette Community Complex March 26th, 7:15 p.m. Silent Auction

2005- 06 Registration for Frères Jacques pre- school Thursday, March 17th 6:00 - 8:00 p.m. Dawson Trail School If you missed it call Denise at 878- 4712

A fifth grader announced at breakfast that, "Today we are having a test on the moon." And her little brother piped up, "Are you gonna let her go? Mom?"

Congratulations

Lorette Busy Bees 4H public speeches were held on Fri., Feb.11. Congratulations to the winners:



Junior speeches: 1st place-Stephanie Walker
2nd place-Ayslynn Clawson

One person Visual-Junior Category: 1st place Hannah Gray
2nd place Justin Rawluk

Intermediate speeches: 1st place-Amy-leigh Gray
2nd place-Serena Ross

Two person visual-Senior category:
1st place Kyra and Karly Gladstone

All the speeches were well done! We would like to thank all who participated! Great job everyone! A Big Thank you to the judges, friends and family who came out for this event.

All these winners were eligible to attend the Steinbach District communications on Sat. March 5, 2005 at Mitchell Middle School.

Second-hand Smoke Hurts Children



Second-hand smoke has serious effects on children's health. Children of parents who smoke"

- ✓ Cough and wheeze more
- ✓ Have more ear infections
- ✓ Go to the hospital more often with bronchitis and pneumonia
- ✓ Have reduced lung function
- ✓ Children with asthma have more attacks.

Second-hand smoke is made up of two kinds of smoke:

Mainstream smoke is the smokes that smokers breath into their lungs and then out into the air.

Sidestream smoke comes from the burning end of the cigarette between puffs.

Second-hand smoke has up to 4,000 chemicals in it. More than 50 of them are known to cause cancer in humans. Three cancer-causing chemical found in second-hand smoke are so dangerous that the North American air quality rules used by most governments say no one should ever be exposed to them.

Sidestream smoke from the burning end of the cigarette has more tar, nicotine, carbon monoxide and other chemicals that cause cancer than the smoke inhaled from the filtered cigarettes by the smoker. When a cigarette burns, about 85% of the smoke that goes into the air is sidestream smoke.

This document was created with Win2PDF available at <http://www.daneprairie.com>.
The unregistered version of Win2PDF is for evaluation or non-commercial use only.